

On Spaces of connected Graphs III

The Ladder Filtration

Jan A. Kneissler

February 1, 2008

Abstract

A new filtration of the spaces of tri-/univalent graphs $\mathcal{B}_m^u$ that occur in the theory of finite-type invariants of knots and 3-manifolds is introduced.

Combining the results of the two preceding articles, the quotients of this filtration are modeled by spaces of graphs with two types of edges and four types of vertices, and an upper bound for $\dim \mathcal{B}_m^u$ in terms of the dimensions of the filtration quotients is given. The degree m up to which $\mathcal{B}_m^u$ is known is raised by two for $u = 0$ and $u = 2$.

0 Definitions and former results

This paper is based heavily on results of the preceding articles [4] and [5]. In this section we briefly repeat the relevant definitions and list results of [4] that will be used.

First recall, that in the context of Vassiliev theory graphs with vertices of order one and three (uni-, trivalent vertices) are subjected to some local relations (i.e. a quotient of a vector space with a basis of such graphs is quotiented by linear combinations of graphs differing only inside a small region in way that is specified by the "relation". Edges that go out off the region are clipped, but the resulting "ends" should not be confused with univalent vertices: all relations in this paper do not involve any univalent vertices. Furthermore, the ends are supposed to be arranged correspondingly in all terms of a relations).

The incoming edges at every trivalent vertex are supposed to be cyclically ordered and the graphs are always drawn such that these orderings correspond with the orientation of the plane. A connected graph with $2m$ vertices will be called *diagram of degree m* .

Definition 0.1

$$\mathcal{B}_m^u := \mathbf{Q}\langle \text{diagrams of degree } m \text{ with } u \text{ univalent vertices} \rangle / (\text{AS}), (\text{IHX})$$

$$\hat{\mathcal{B}}_m^u := \mathbf{Q}\langle \text{diagrams of degree } m \text{ with } u \text{ univalent vertices} \rangle / (\text{AS}), (\text{IHX}), (\text{x})$$

where (AS), (IHX) and (x) are the following local relations ((x) is rather a family of relations):

$$\begin{array}{c} \diagup \quad \diagdown \\ \quad \quad \quad \end{array} = - \begin{array}{c} \diagup \quad \diagdown \\ \quad \quad \diagup \quad \diagdown \end{array} \quad (\text{AS})$$

$$\begin{array}{c} | \\ | \\ | \end{array} - \begin{array}{c} | \quad | \\ | \quad | \end{array} - \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} = 0 \quad (\text{IHX})$$

$$\begin{array}{c} \text{---} \\ | \\ | \\ | \end{array} = 0 \quad (\text{x})$$

In [4] we have found a set of consequences of the relations in $\hat{\mathcal{B}}_m^u$ that will be repeated here, for the readers convenience:

$$\begin{array}{c} \text{---} \diagup \text{---} \diagdown \\ \text{---} \diagdown \text{---} \diagup \end{array} = \begin{array}{c} \text{---} \diagup \text{---} \diagdown \\ \text{---} \diagdown \text{---} \diagup \end{array} + \begin{array}{c} \text{---} \diagup \text{---} \diagdown \\ \text{---} \diagdown \text{---} \diagup \end{array} = - \begin{array}{c} \text{---} \diagup \text{---} \diagdown \\ \text{---} \diagdown \text{---} \diagup \end{array} = 0 \quad (\text{t})$$

$$\begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} + \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} \quad (\text{s})$$

$$\begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} \quad (\text{LS})$$

$$\begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} + \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} + \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = 0 \quad (\text{LIHX})$$

$$2 \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} + \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = 0 \quad (\text{LI})$$

$$4 \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} = \begin{array}{c} \diagup \text{---} \diagdown \\ \diagdown \text{---} \diagup \end{array} \quad (\text{LL})$$

The last four relations involve subgraphs of a special form, that will be called *ladders*. A ladder with $n \geq 2$ rungs is called n -ladder; it consists of $3n + 2$ edges. The four edges connecting the ladder with the rest of a diagram are called *ends* of the ladder. A 2-ladder is also called *square*. An n -ladder is called *odd* and *even* according to the parity of n . The set of all ladders of a given diagram is partially ordered by the subgraph relation. Ladders that are maximal elements with respect to this relation are called *maximal* ladders.

Each of the last four equations, involving ladders, represents a whole family of relations according to the following rules:

1. Each of the example ladders in the pictures above may be replaced by a n -ladder of same parity ($n \geq 2$).
2. The relation must remain homogeneous (i.e. each term of a relation must get the same total number of rungs).

The main result of [4] (Theorem 3, "square-tunneling relation") shall also be restated:

Theorem 0 *Moving two rungs from a p -ladder to another q -ladder inside a diagram (thus the new diagram has a $(p-2)$ - and a $(q+2)$ -ladder) does not change its value in $\hat{\mathcal{B}}_m^u$ if $p \geq 4$, $q \geq 2$.*

1 Introduction

In the first of this series of articles we mainly examined the spaces $\hat{\mathcal{B}}_m^u$. The second article dealt with a sub-algebra Λ_0 of the algebra Λ that has been defined in [8]. Nevertheless not $\hat{\mathcal{B}}_m^u$ and Λ_0 but rather $\mathcal{B}_m^u$ and Λ are the spaces that are of significance for the theory of finite type invariants of knots and of 3-manifolds, respectively.

So we made life too easy in [4] by using the additional relation (x) and in [5] we possibly missed some part of Λ since only the elements x_n were considered.

But recall that Λ acts on $\mathcal{B}_m^u$ and that Λ is isomorphic (as graded vector space) to $\bigcup_{m \geq 1} \mathcal{B}_m^0$. So on one side (in [4]) we ignored the elements of Λ_0 (of degree ≥ 1 at least), whereas on the other side (in [5]) we had a somehow “orthogonal” point of view.

Thus it is quite natural to look for a synthesis of both approaches that “reassembles” $\mathcal{B}_m^u$ out of the fragments $\hat{\mathcal{B}}_m^u$ and Λ_0 .

1.1 The ladder filtration

The statement of Theorem 0 can be reformulated as follows: A square that is part of a n -ladder with $n \geq 4$ is called *free square* (since it may move around freely across ladders). The result of removing a free square (i.e. reducing the number of rungs of a ladder by 2) is called a *reduction*. For each diagram one obtains after a finite number of reductions a unique diagram, which will be called *complete reduction*. Now Theorem 0 implies that diagrams of same degree and same complete reduction represent identical elements of $\hat{\mathcal{B}}_m^u$. In other words, diagrams may be viewed as combination of completely reduced diagrams (containing ladders of length 2 or 3 only) together with a “cloud” of free squares.

We suggest to filter $\hat{\mathcal{B}}_m^u$ by the number of such free squares and the number of odd and even ladders, which motivates the following Definitions 1.1 and 1.3.

Definition 1.1 For a diagram D and $i \geq 2$ let $n_i(D)$ denote the number of maximal ladders of length i in D . Let $\delta(D)$ be the triple of integers counting the number of free squares, odd ladders and even ladders:

$$\delta(D) := \left(\sum_{i \geq 2} n_i(D) \cdot \left\lfloor \frac{i-2}{2} \right\rfloor, \sum_{i \geq 1} n_{2i+1}(D), \sum_{i \geq 1} n_{2i}(D) \right)$$

Proposition 1.2 Let D denote a diagram of $\hat{\mathcal{B}}_m^u$ with $\delta(D) = (f, o, e)$, then

$$\begin{aligned} f, o, e &\geq 0 \\ o + e = 0 &\Rightarrow f = 0 \\ 4f + 6o + 4e &\leq 2m - u \\ 2f + 2o + e &\leq m - u + 1. \end{aligned}$$

The first two conditions are obvious, the third states that the numbers of vertices in ladders does not exceed the total number of trivalent vertices in D and the forth is obtained by the fact that the cyclomatic number of D is $m - u + 1$ and that the ladders in D contain $\sum n(i) \cdot (i - 1)$ linearly independent cycles.

Definition 1.3 Let $T(m, u)$ denote the subset of $\mathbf{Z}^3$ consisting of all triples (f, o, e) that satisfy the four conditions of Proposition 1.2. Let $\prec$ denote the ordering on $T(m, u)$ that is induced by the lexicographical order of $\mathbf{Z}^3$. Let $\mathcal{F}_{f,o,e} \hat{\mathcal{B}}_m^u$ denote the subspace of $\hat{\mathcal{B}}_m^u$ that is spanned by all diagrams D with $(f, o, e) \preceq \delta(D)$.

We get a decreasing filtration (t is the maximal element of $T(m, u)$):

$$\hat{\mathcal{B}}_m^u = \mathcal{F}_{0,0,0} \hat{\mathcal{B}}_m^u \supset \mathcal{F}_{0,0,1} \hat{\mathcal{B}}_m^u \supset \dots \supset \mathcal{F}_t \hat{\mathcal{B}}_m^u.$$

The aim of this article is to obtain some information about the filtration quotients that will be denoted $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$. A strong source for the hope that this filtration is helpful for the study of $\hat{\mathcal{B}}_m^u$ (and thus $\mathcal{B}_m^u$) is the observation that the quotients probably are independent of f (up to a shift of indices). We have found evidence for this in many situations, which encouraged us to formulate it as a conjecture.

Conjecture 1.4 There are isomorphisms $\mathcal{G}_{f_1,o,e} \hat{\mathcal{B}}_{m+2f_1}^u \cong \mathcal{G}_{f_2,o,e} \hat{\mathcal{B}}_{m+2f_2}^u$ for all integers m, u, f_1, f_2, o, e satisfying $(f_i, o, e) \in T(m + 2f_i, u)$.

1.2 Feynman graphs


To represent the elements of $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$, we have to specify the places where odd and even ladders are situated. Therefore, we will introduce graphs that contain vertices of degree 4 and a second type of edges. The similarity to graphs used in the physics of elementary particles is so striking (but meaningless) that we could not resist calling them Feynman graphs. Some authors already name the diagrams of $\mathcal{B}_m^u$ this way, (these are also Feynman graphs in our sense, so our terminology extends the old one). Nevertheless, to avoid confusion, we will always say “diagram” when are talking about elements of $\mathcal{B}_m^u$, $\hat{\mathcal{B}}_m^u$ or $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ and “Feynman graph” when we are speaking about the spaces $\mathcal{C}$ that will be defined in this subsection.

Definition 1.5 A *Feynman graph* is a connected multi-graph with two types of edges called *normal edges* and *photon edges*. The vertices all have valency 1, 3 or 4. The ends of photon edges must be trivalent and no more than one photon edge may arrive at a trivalent vertex. The ends of a photon edge are called *photon vertices* and the other trivalent vertices are called *normal vertices*. At every normal vertex, a cyclic ordering of the three incoming edges is specified, but at the other vertices, the edges are not ordered in any way.

In pictures, the edges are drawn with straight — and wavy ~ lines. The cyclic ordering at normal edges is assumed counter-clockwise in all pictures. The restrictions imply that Feynman graphs may only contain the following four types of vertices:



The *degree* of a Feynman graph is a quadruple of integers (m, u, p, q) where u denotes the number of univalent vertices, p the number of photon edges, q the number of tetravalent vertices and $2m - u - 6p - 4q$ is equal to the number of normal vertices (m is always an integer). Here are two examples of Feynman graphs of degree $(8, 4, 1, 1)$:



Finally, let $\mathcal{C}_{m,u,p,q}$ denote the $\mathbf{Q}$ -vector space that is spanned by all Feynman graphs of degree (m, u, p, q) . The reader might anticipate that the preceding definitions will ensure the spaces $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ are quotients of these spaces of Feynman graphs. In fact, there is an epimorphism $\varphi_f : \mathcal{C}_{m,u,o,e} \rightarrow \mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{m+2f}^u$ for all allowed m, u, f, o, e (for simplicity, we denote the all maps with different parameters m, u, o, e by the same symbol φ_f , and hope this will not lead to confusion).

The big question is, how do the kernels of the maps φ_f look like? The best way to answer this question would be to give a complete list of local relations that generate the kernels for all possible parameters m, u, f, o, e . But we do not know if such an answer is possible. Furthermore, even if there exists such a list, it still might be infinite and almost impossible to find.

At the time, we are only able to present some of the relations we could make out (see Theorem 1). They all turn out to be independent of the number f , which strengthens our belief in Conjecture 1.4, which states $\ker \varphi_{f_1} = \ker \varphi_{f_2}$.

2 Results

It is not hard to guess how the maps φ_f are constructed; for any Feynman graph G let $r(G)$ denote the diagram that is obtained by making the following two substitutions at every tetravalent vertex

and every photon edge of G , respectively:

$$\begin{array}{ccc} \times & \rightarrow & \square \\ \text{wavy line} & \rightarrow & \square \end{array}$$

Furthermore, for a diagram D of $\hat{\mathcal{B}}_m^u$ containing at least one ladder, let $s(D)$ denote the diagram that is obtained by adding a square to an arbitrarily chosen ladder of D :

$$\square \rightarrow \square \square$$

If there are two or more ladders in D then $s(D)$ is not uniquely defined by this, but by the square-tunneling relation (Theorem 0) ensures that different choices yield the same element in $\hat{\mathcal{B}}_{m+2}^u$.

Theorem 1 *The map $s^f \circ r$ from Feynman graphs to diagrams extends to a well-defined epimorphism $\varphi_f : \mathcal{C}_{m,u,o,e} \rightarrow \mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{m+2f}^u$ for all $(f, o, e) \in T(m+2f, u)$. Elements of the subsequent forms are annihilated by these epimorphisms:*

- linear combinations that look locally like one of the following:

$$\text{Y-vertex} + \text{X-vertex with loop} , \quad (\text{AS})$$

$$\text{I} - \text{H} - \text{X} , \quad (\text{IHX})$$

$$\text{I} + \text{wavy line} + \text{X} , \quad (\text{LIHX})$$

$$\text{X} + \text{X} - \text{X} - \text{X} , \quad (4\text{T}_a)$$

$$\text{wavy line} + \text{wavy line} - \text{wavy line} - \text{wavy line} , \quad (4\text{T}_b)$$

- graphs that contain a cycle of length 1, 2 (if $m > 2$) or 3,
- graphs that contain a cycle of length 4 of one of the following types:

$$\begin{array}{cccccc} \square & , & \square & , & \square & , & \square & , & \square \\ \square & , & \square & , & \square & , & \square & , & \square \end{array}$$

- graphs that contain a cycle of length 5 of one of the following types:

$$\text{pentagon} , \text{pentagon} , \text{pentagon} .$$

The fact that there are a lot of forbidden cycles lets one expect that – at least in low degrees – the spaces $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ have low dimensions. For instance, using the Euler characteristics of compact orientable surfaces, it is not hard to show that a Feynman graph with genus g , q tetravalent vertices and $u < q + 6(1 - g)$ univalent vertices contains a cycle of length ≤ 5 . This implies for $u < 6$ that $\mathcal{G}_{0,0,0} \hat{\mathcal{B}}_m^u$ is spanned by non-planar graphs. In [2] it has been conjectured that $\mathcal{B}_m^0$ is spanned by planar diagrams; if this was true, then $\mathcal{G}_{0,0,0} \hat{\mathcal{B}}_m^0 = 0$ for all m .

2.1 Upper bounds for $\dim \mathcal{B}_m^u$

We now focus on the spaces $\mathcal{B}_m^u$ rather than $\hat{\mathcal{B}}_m^u$. It is clear that information about the filtration quotients $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ will somehow provide upper bounds for $\dim \mathcal{B}_m^u$. With the next statement, we will make this more explicit.

Theorem 2 *Assume that we are given upper bounds for the dimensions of $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ that are independent of f , i.e. there exist numbers $\mu_{m,u,o,e}$ satisfying $\dim \mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{m+2f}^u \leq \mu_{m,u,o,e}$ for all f . With the abbreviations $q_n := \lfloor \frac{n^2}{12} + \frac{1}{2} \rfloor$ and $\delta_{i,j} := \begin{cases} 1 & \text{for } i=j \\ 0 & \text{for } i \neq j \end{cases}$, the following inequality then holds for all $m \geq u \geq 0$:*

$$\dim \mathcal{B}_m^u \leq \sum_{j=u}^m \left((1 + q_{m-j}) \mu_{j,u,0,0} + q_{m-j+3} \sum_{o+e \geq 1} \mu_{j,u,o,e} \right) + q_{m-6} \cdot \delta_{u,4}.$$

The inner sum runs over all $o, e \geq 0$ with $o + e \geq 1$, $6o + 4e \leq 2m - u$ and $2o + e \leq m - u + 1$.



The idea, of course, is to obtain such upper bounds by use of Theorem 1, by simply taking

$$\mu_{m,u,o,e} := \dim \left(\mathcal{C}_{m,u,o,e} \text{ / relations listed in Theorem 1} \right).$$

2.2 Experimental data


The spaces $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ have been calculated for $m \leq 12$ and $2 \leq u \leq 6$. This has been done by establishing upper and lower bounds for $\dim \mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$. The computation of upper bounds is based on Theorem 1; lower bounds are derived by comparing the statement of Theorem 2 with the known exact values of $\dim \mathcal{B}_m^u$ of [3].

As predicted by Conjecture 1.4, the calculated filtration spaces do not depend on f (up to the shift in the degree m). All but 15 of the spaces in question are trivial. In the non-trivial cases, $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$ has dimension one. In the following table we list these spaces together with a Feynman graph that is mapped by φ_f to a generator (the case $u = 0$ which can be derived from $u = 2$ is also shown). The non-triviality of these diagrams (as elements of $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^u$) has been verified using weight systems that come from the Lie-algebras $\mathfrak{gl}$ and $\mathfrak{so}$.

$\mathcal{G}_{0,0,0} \hat{\mathcal{B}}_1^0$	$\mathcal{G}_{0,0,0} \hat{\mathcal{B}}_2^2$	$\mathcal{G}_{f,0,1} \hat{\mathcal{B}}_{4+2f}^4$ ($0 \leq f \leq 4$)	$\mathcal{G}_{0,0,0} \hat{\mathcal{B}}_6^6$	$\mathcal{G}_{f,0,2} \hat{\mathcal{B}}_{7+2f}^6$ ($0 \leq f \leq 2$)	$\mathcal{G}_{f,1,0} \hat{\mathcal{B}}_{8+2f}^6$ ($0 \leq f \leq 2$)	$\mathcal{G}_{f,2,0} \hat{\mathcal{B}}_{10+2f}^6$ ($0 \leq f \leq 1$)
						

Remark 2.1 The upper bounds for $\dim \mathcal{B}_m^u$ of Theorem 2 are sharp for $m \leq 12$.

Considering the results of [3], one might conjecture that $\hat{\mathcal{B}}_m^u$ is spanned by diagrams the following type (i.e. caterpillars with an even number of legs on each of its body segments):



This conjecture is quite daring and we have the feeling that it will turn out false. Nevertheless let us mention that it implies the following weaker statement, that is more likely to be true:

Conjecture 2.2 *For $m > u + 1 > 0$ and $m > 5 \left(\frac{u}{2} - 1 \right)$ the filtration quotients $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{m+2f}^u$ are trivial for all possible parameters f, o, e .*

If Conjecture 2.2 is true then, due to Theorem 2, $\dim \mathcal{B}_m^u$ grows at most quadratically with m for fixed u . To be more specific: together with Theorem 2 and taking into account the data of the table above, Conjecture 2.2 implies

Conjecture 2.3

$$\begin{aligned} \dim \mathcal{B}_{m-1}^0 &= \dim \mathcal{B}_m^2 \leq \left\lfloor \frac{(m-2)^2}{12} + \frac{3}{2} \right\rfloor \\ \dim \mathcal{B}_m^4 &\leq \left\lfloor \frac{(2m-7)^2}{24} + 1 \right\rfloor \\ \dim \mathcal{B}_m^6 &\leq \left\lfloor \frac{(2m-11)^2}{12} + \frac{3}{2} \right\rfloor. \end{aligned}$$

These conjectural inequalities are fulfilled and even sharp for $m \leq 12$.

Remark 2.4 The restriction of Conjecture 2.2 to the case $u = 0$ (or equivalently $u = 2$) is equivalent to Conjecture 1.2 of [5] (claiming $\Lambda = \Lambda_0$, i.e. Vogel's algebra Λ is generated by the elements x_n with $n \geq 1$ already, see also [8]). It is known that $\Lambda = \Lambda_0$ for degrees ≤ 12 due to the following

Computational result: $\dim \mathcal{G}_{f,o,e} \hat{\mathcal{B}}_m^2 = 0$ for $3 \leq m \leq 14$.

This also implies that the first inequality of Conjecture 2.3 is correct and sharp up to $m = 14$. The algorithm of the computer program that checked triviality for the previously unknown degrees $m = 13$ and $m = 14$ is based on the relations of Theorem 1 and the fact that in all considered degrees it suffices to take into account only Feynman graphs of a very special form (graphs in which there is a path going through all non-normal vertices and through all photon edges, not containing any normal vertices).

Remark 2.5 In chapter 7 of [6], using results of Petersen and Chvátal, $\dim \mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{13}^2 = 0$ is shown "by hand", i.e. without aid of a computer program. Petersen's theorem states that any $(2n)$ -regular graph has a 2-factor; Chvátal's theorem gives a sufficient condition for a graph being path-hamiltonian in terms of its degree sequence.

3 Proof of Theorem 1

The map r replaces tetravalent vertices by squares and photon edges by ladders of length 3. Because of relation (s), the result does not depend (in $\hat{\mathcal{B}}_m^u$) on the different possible choices of permutations of the four incoming edges.

Due to the square-tunneling relation (Theorem 0), adding squares to existing ladders is a well-defined operation in $\hat{\mathcal{B}}_m^u$. Furthermore, the degree of Feynman graphs has been chosen to ensure that r is degree-preserving with respect to the m - and u -components of the degree and s increases the total degree m by 2. So $s^f \circ r$ extends to a homomorphism from $\mathcal{C}_{m,u,o,e}$ to $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{m+2f}^u$ (there is no problem with the fact that s is only defined for diagrams with ladders, because $o = e = 0$ implies $f = 0$, i.e. s is never applied on diagrams without ladders).

Obviously, every diagram of $\mathcal{G}_{f,o,e} \hat{\mathcal{B}}_{m+2f}^u$ is the image of some Feynman graph in $\mathcal{C}_{m,u,o,e}$ (to find the Feynman graph, simply take the complete reduction of the diagram and invert r).


So the only thing that remains to prove Theorem 1 is to verify the list of relations. The relations (AS), (IHx), (LIHX) are mapped by φ_f to the relation in $\hat{\mathcal{B}}_{m+2f}^u$ with the same names; $(4T_a)$ and $(4T_b)$ are special cases of the fact that, taking the (consistently signed) sum of all possible ways of attaching an edge at all entries of a region without univalent vertices, one obtains a trivial element of $\mathcal{B}_m^u$, which is obviously also trivial in $\hat{\mathcal{B}}_m^u$ (see Lemma 3.1 a) of [4] for more details).

3.1 Loops and double edges

There are three possible situations with cycles of length 1:



The first is trivial by (AS), the last two are trivial because of the relation (x). Next, we consider cycles of length 2 (i.e. double edges):




In the first case, if there is at least one additional trivalent vertex in the diagram, we may apply relation (x). Only in the cases $m = 1, u = 0$ and $m = 2, u = 2$, there exist diagrams having a cycle of the first type, but no other trivalent vertices (see the first two columns of the table on the preceding page). This explains why we have to make the requirement $m > 2$.

The cases 2, 3 are killed by relation (t) and cases 4, 5, 6 are mapped by φ_f to diagrams with more than f free squares, because the length of two ladders are added. In the last case we may apply relation (x).

3.2 The flying-trick


In many situations we apply a useful procedure that we call *flying*. It can generally be described like this: take the cycle out of the plane, make a half twist around an appropriate axis and put it back into the plane. By (AS) each normal vertex causes a change of sign. Then use relations (AS), (IHX), (4T_a), (4T_b) and (LIHX) to take the cycle into its original form, modulo simpler cycles. If the result has opposite sign, then the described operation yields an equation that allows to express the cycle in terms of simpler diagrams.

We illustrate this trick for two examples of triangles that in this way can be expressed by diagrams with double edges (relations (LIHX) and (4T_a) are used):




3.3 Triangles

There are thirteen possible cycles of length 3; let us first discuss the easy ones:



The first is killed by relation (t), the second by (x). In the third and forth situations, the length of the ladder may be increased by including the edge between the two normal vertices. In the last five cases one may successfully apply the flying-trick. Now we have to take a look at the tricky triangles:



Let us assume that all free squares are inserted at the ladder on the left side, when φ_f is applied to these graphs. Then the ladders in the upper and right corner will have length 2 or 3. Here we need a result of [4] (equation (37) in section 4.4), which states the following local relation:

$$\begin{aligned}
 \text{Diagram 1} &= \text{Diagram 2} - \text{Diagram 3} + \text{Diagram 4} \\
 &\quad - \text{Diagram 5} + \text{Diagram 6}
 \end{aligned}
 \tag{*}$$

The diagrams with rounded boxes shall be interpreted as linear combination of 4 diagrams by expanding each box the following manner:

$$\text{Diagram with rounded box} = \frac{1}{2} \text{Diagram 1} + \frac{1}{2} \text{Diagram 2}$$

We may apply (*) in all four cases and obtain diagrams in which the triple (number free squares, number of odd ladders, number of even ladders) is lexicographically greater than in the original diagram.

3.4 Simple cycles of length 4 and 5

Again we start with the simple cases:

$$\text{Diagram 1}, \text{Diagram 2}, \text{Diagram 3}, \text{Diagram 4}, \text{Diagram 5} .$$

The first contains a square that is not coming from a tetravalent vertex, the next three can be handled with the flying-trick and the last is due to the relation (LL).

There are two 5-cycles that can be reduced by the flying-trick to the first and last of the 4-cycles above:

$$\text{Diagram 6}, \text{Diagram 7} .$$

3.5 Pentagons with a photon edge

Perhaps the most surprising (and hardest to prove) statement is that cycles of this type (i.e. containing a photon edge and three normal vertices) are annihilated by φ_f :

$$\text{Diagram 8} .$$

To show this, we start with the observation that the occurrence of one of the following four situations allows to produce a ladder of length 4:

$$\text{a) Diagram 9}, \quad \text{b) Diagram 10}, \quad \text{c) Diagram 11}, \quad \text{d) Diagram 12} .$$

In case a) and c) this is done by the flying trick. Case b) corresponds to relation (LI) and d) can be reduced to a trivial diagram (because of relation (t)) and type c) and b) in the following way:

$$2 \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}$$

Let us recall a result of [4] (equation (38), section 4.4, it involves a parameter $n \geq 1$), which looks for $n = 1$ like this:

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}$$

If we glue the upper left ends together like this $\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \rightarrow \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}$, we obtain

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \equiv - \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}.$$

By the sign $\equiv$, we mean “modulo terms containing a 4-ladder”. This congruence is true, because three of the diagrams in the middle are of type a), b) and d). On the other hand, if we use (IHX) and neglect another diagram of type d), we get

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \equiv - \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}.$$

One may repeat this argument for arbitrary odd ladders instead of the 3-ladder (using the general version of equation (38) of [4]). Thus we have shown that the following two Feynman graphs are mapped to equivalent diagrams in the corresponding filtration quotient:

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}$$

This finally leads to the following (oval boxes to be interpreted as in section 3.3):

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \frac{1}{2} \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \frac{1}{2} \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \\ = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \frac{1}{2} \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = -\frac{1}{2} \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} + \frac{1}{2} \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}.$$

As we have seen before, these two 4-cycles lie in the kernel of φ_f .

3.6 Cycles with a photon edge and non-normal vertices

In the preceding subsection we have seen that the occurrence of a pentagon with a photon edge allows to increase the number of free squares. This immediately applies in the following two situations (we have to sacrifice ladders when reducing the pentagon, but this is acceptable since we gain free squares):

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}, \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}.$$

The only three remaining cycles in our list are the following:

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}, \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}, \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}.$$

They all can be reduced by the flying-trick, if we realize that

$$\begin{array}{c} \text{Diagram 1} \end{array} = \begin{array}{c} \text{Diagram 2} \end{array} + \left(\text{terms containing } \begin{array}{c} \text{Diagram 3} \end{array} \right)$$

and similar equations with one or both of the squares replaced by 3-ladders. $\square$

4 Proof of Theorem 2

Recall that $\mathcal{B}_m^u$ may be viewed as Λ_0 -module (see [5] for a definition of Λ_0 and its elements x_n). $\hat{\mathcal{B}}_m^u$ is obtained by quotienting $\mathcal{B}_m^u$ by all elements that are multiples of Λ_0 -elements of degree ≥ 1 . So by Theorem 1, $\mathcal{B}_m^u$ is spanned the following set: ($\mathcal{G}_n \Lambda_0$ denotes the degree n part of Λ_0)

$$\{ \lambda \varphi_f(d) \mid \lambda \in \mathcal{G}_i \Lambda_0, \ d \in \mathcal{C}_{m-2f-i,u,p,q}, \ (f,p,q) \in T(m-i,u) \}.$$

In other words: we obtain a spanning set for $\mathcal{B}_m^u$ by taking basis elements of $\mathcal{C}_{m',u,p,q}$ with $m' \leq m$ and blowing up their degree to the correct value by inserting free squares and elements of Λ_0 in all possible ways.

To get the desired inequality, we still have to reduce the number of generators a little bit. We therefor have to use the fact that $\mathcal{B}_m^u$ is not a free Λ_0 -module, what has been discovered in [3].

By Theorem 0 we know that the following local relation holds in $\hat{\mathcal{B}}_m^u$:

$$\begin{array}{c} \text{Diagram 1} \end{array} = \begin{array}{c} \text{Diagram 2} \end{array}$$

But in view of Corollary 2.4 of [4], one can hope for more: there must also exist a similar local relation in $\mathcal{B}_m^u$, involving elements of Λ_0 (a local relation is, in the terminology of [4], nothing else than a trivial linear combination in $F(6)$; Corollary 2.4 of [4] gives such a trivial element in $\hat{F}(6)$, the Λ_0 -quotient of $F(6)$). The corresponding identity in $F(6)$ in fact looks like this:

$$\begin{aligned} 3 \begin{array}{c} \text{Diagram 1} \end{array} &= 3 \begin{array}{c} \text{Diagram 2} \end{array} + 2x_3 \begin{array}{c} \text{Diagram 3} \end{array} - 2x_3 \begin{array}{c} \text{Diagram 4} \end{array} \\ &+ x_3 \begin{array}{c} \text{Diagram 5} \end{array} + t(\dots) \end{aligned}$$

Closing this relation in the following way $\begin{array}{c} \text{Diagram 6} \end{array} \rightarrow \begin{array}{c} \text{Diagram 7} \end{array}$, we obtain for $n \geq 3$:

$$x_n \begin{array}{c} \text{Diagram 8} \end{array} = (x_{n+2} + \frac{2}{3}x_3x_{n-1}) \begin{array}{c} \text{Diagram 9} \end{array} - \frac{2}{3}x_3x_n \begin{array}{c} \text{Diagram 10} \end{array} + \frac{1}{3}x_3x_n \begin{array}{c} \text{Diagram 11} \end{array} + t(\dots)$$

This means that an element of the form $t^i x_n \varphi_f(d)$ with $n \geq 3$ and $f > 0$ can be expressed in terms of elements containing more t -s or a Λ_0 -factor of higher degree. There is one exception: if $d = \mathbf{X} \in \mathcal{C}_{4,4,0,1}$ and $f = 1$, then there is no trivalent vertex in $\varphi_1(d)$ other than the eight vertices of the ladder. We thus may not apply the upper relation to this case.

In all other cases we may lower the number of free squares successively, as long as there is an x_n with $n \geq 3$ present. Consequently, $\mathcal{B}_m^u$ is spanned by elements of the three types (the third only applies to $u = 4$):

- a) $\lambda \varphi_0(d)$ with $\lambda \in \mathcal{G}_i \Lambda_0$ and $d \in D_{m-i,u,p,q}$, $(0,p,q) \in T(m-i,u)$,
- b) $t^i \varphi_f(d)$ with $f \geq 1$ and $d \in D_{m-i-2f,u,p,q}$, $(f,p,q) \in T(m-i,u)$,
- c) $\lambda \varphi_1(\mathbf{X})$ with $\lambda \in \mathcal{G}_{m-6} \Lambda_0$.

To make this argument more convincing, we would have to define a filtration of $\mathcal{B}_m^u$ according to the lexicographical ordering of the pairs (number of t -s, sum of degrees of x_n -s with $n \geq 3$), but we omit this technical complication.

By Corollary 1.6 of [5] we know that $\dim \mathcal{G}_n \Lambda_0 \leq 1 + q_n$. So the number of linearly independent elements of type a) is at most

$$\sum_{i=m-u}^m \sum_{p,q} (1 + q_i) \dim D_{m-i,u,p,q}.$$

The outer sum starts with $i = m - u$, because $D_{m,u,p,q}$ is always trivial if $m < u$. The inner sum runs over all allowed p, q i.e. all p, q that satisfy $(0, p, q) \in T(m - i, u)$.

The number of independent elements of type b) is

$$\sum_{j=u}^m \sum_{p+q \geq 1} \left\lfloor \frac{m-j}{2} \right\rfloor \dim D_{j,u,p,q}$$

because $\lfloor \frac{m-j}{2} \rfloor$ is the number of pairs (i, f) satisfying $i \geq 0, f \geq 1, j + i + 2f = m$. The inner sum runs over all allowed p, q with $p + q \geq 1$ (we can only add free squares if there is at least one ladder).

There are $1 + q_{m-6}$ elements of type c), but $\lambda = t^{m-6}$ has already been counted in b). So the additional contribution to the dimension by type c) elements is at most $q_{m-j} \delta_{u,4}$.

Substituting $j = m - i$ and using $1 + q_n + \lfloor \frac{n}{2} \rfloor = q_{n+3}$, all three contributions sum up to the inequality given in Theorem 2. $\square$

References

- [1] D. Bar-Natan, *On the Vassiliev knot invariants*, Topology **34** (1995), 423-472.
- [2] S. V. Chmutov, S. V. Duzhin, A. I. Kaishev, *The algebra of 3-graphs*, Program System Institute (Pereslavl-Zalessky, Russia) preprint, December 1997.
- [3] J. A. Kneissler, *The number of primitive Vassiliev invariants up to degree twelve*, e-print archive (<http://xxx.lanl.gov>), **q-alg/9706022**.
- [4] J. A. Kneissler, *On spaces of connected graphs I: Properties of ladders*, Proc. Internat. Conf. "Knots in Hellas '98", Series on Knots and Everything, vol. **24** (2000), 252-273.
- [5] J. A. Kneissler, *On spaces of connected graphs II: Relations in the algebra Λ* , Jour. of Knot Theory and its Ramif. Vol. **10**, No. 5 (2001), 667-674.
- [6] J. A. Kneissler, *Die Kombinatorik der Diagrammalgebren von Invarianten endlichen Typs*, doctoral thesis (Rheinische Friedrich-Whilhelms-Universität Bonn), Bonner Mathematische Schriften **325** (1999).
- [7] V. A. Vassiliev, *Cohomology of knot spaces*, Theory of Singularities and its Applications (ed. V. I. Arnold), Advances in Soviet Math., **1** (1990) 23-69.
- [8] Pierre Vogel, *Algebraic structures on modules of diagrams*, Université Paris VII preprint, July 1995 (revised 1997).

e-mail: jan@kneissler.info
<http://www.kneissler.info>